

Übungen zur Theorie der Kondensierten Materie II SS 16

PROF. J. SCHMALIAN

Blatt 2

M. S. SCHEURER, B. JEEVANESAN

Besprechung 06.05.2016**1. Dyson equation**

(6 Points)

We want to describe a particle on a lattice that interacts with an impurity at the origin. The sites of the lattice are numbered by an index i . Thus the state $|i\rangle$ refers to a position eigenstate located on site i .

Consider a Hamiltonian H that is the sum of two parts

$$H = H_0 + V = \sum_{ij} H_{ij} |i\rangle\langle j| + V_0 |0\rangle\langle 0|.$$

Assume that we know everything about the problem involving the Hamiltonian H_0 . We now want to study the effect of an impurity that sits at site 0. This is represented by the second term. The full single-particle Green's function is a matrix given by

$$G^{-1} = \omega - H_0 - V.$$

We define the bare Green's function G_0 as

$$G_0^{-1} = \omega - H_0.$$

Show that

$$G = \frac{1}{1 - G_0 V} G_0$$

holds. Now expand the fraction as a geometric series and compute the matrix element $\langle i|G|j\rangle$. You should find

$$\langle i|G|j\rangle = \langle i|G_0|j\rangle + V_0 \frac{\langle i|G_0|0\rangle\langle 0|G_0|j\rangle}{1 - V_0\langle 0|G_0|0\rangle}.$$

2. Asymptotic behavior of Green's functions

(10 Points)

In this problem we will analyze the large- ω behavior of the Green's functions $G_{AB}^r(\omega)$, $G_{AB}^a(\omega)$ and $G_{AB}^c(\omega)$ for fermions $A = B^\dagger$.

- (a) The Lehmann representation for $G^r(\omega)$ was derived in the lectures. Find a similar representation for $G^a(\omega)$ and $G^c(\omega)$ following the arguments used for $G^r(\omega)$. Schematically you will find representations of the form

$$\begin{aligned} G^r(\omega) &= \sum_{lm} \frac{N_r(l, m)}{\omega + E_l - E_m + i0^+} \\ G^a(\omega) &= \sum_{lm} \frac{N_a(l, m)}{\omega + E_l - E_m - i0^+} \\ G^c(\omega) &= \sum_{lm} \frac{N_1(l, m)}{\omega + E_l - E_m + i0^+} + \frac{N_2(l, m)}{\omega + E_l - E_m - i0^+}, \end{aligned}$$

for some appropriate N_r, N_a, N_1 and N_2 that you should determine. (5 Pts.)

(b) Now use the fermionic anti-commutation relations to show that

$$\sum_{l,m} N_r(l, m) = \sum_{l,m} N_a(l, m) = \sum_{l,m} [N_1(l, m) + N_2(l, m)] = 1$$

Use this to determine the asymptotic behavior of the Green's functions for $\omega \rightarrow \infty$.
(5 Pts.)